Hip Luxation in Dogs

What is Hip Luxation?

Hip luxation, also known as a dislocated hip, occurs when the head of the femur (thigh bone) pops out of the hip socket. This usually happens due to trauma or, in some cases, due to severe hip dysplasia, a genetic condition affecting hip joint stability. Both types of luxation cause pain and difficulty walking, but they have different underlying causes.

Causes of Hip Luxation

1.Traumatic Hip Luxation

-Cause: Typically caused by a specific injury, such as being hit by a car, a fall, or an unexpected twist while running.

-Features: The hip joint is usually otherwise healthy, and the main problem is the dislocation itself.

-Treatment: Closed reduction (manual repositioning) or open reduction (surgical repositioning) often works well to restore normal joint function. Recovery is typically good with proper care.

2.Dysplastic Hip Luxation

-Cause: Caused by hip dysplasia, a genetic condition where the hip joint doesn't form properly during growth, leading to instability. Dogs with hip dysplasia often have shallow hip sockets that make the joint prone to dislocating.

-Features: The hip is chronically unstable due to its abnormal shape, which means luxation can occur even with minor movements or no trauma at all.

-Treatment: Dogs with dysplastic hip luxation may benefit from different surgical options, like Femoral Head and Neck Osteotomy (FHNO) or Total Hip Replacement (THR), which help manage both pain and joint instability in the long term.

Signs of Hip Luxation

- Limping or holding one leg up
- Pain, especially when the leg is moved
- Swelling or stiffness around the hip
- Change in posture, with the leg held outward or at an unusual angle

Diagnosis

Your vet will perform a physical examination and likely recommend imaging, such as X-rays, to determine the type of luxation and if there are any underlying issues, such as hip dysplasia.

Treatment Options

Treatment varies based on the type of luxation, the severity of the injury, and whether other structures around the hip joint are damaged:

1. Closed Reduction (Non-Surgical)

This is a non-surgical method where the vet manually pushes the femur back into place. This works best if done soon after the injury (ideally within 72 hours). To keep the hip in place while it heals, your dog may need a special sling or bandage for a couple of weeks.

- Success Rate: Around 50%, especially if done promptly after the injury.
- Risks: The hip may dislocate again, especially if there's ligament damage.

2. Open Reduction (Surgical)

If closed reduction isn't successful or if the dislocation is severe, surgery may be necessary. In an open reduction, the vet will surgically reposition the femur and secure it in place using sutures, pins, or other techniques. This helps stabilize the hip joint and prevent it from dislocating again.

- Success Rate: Approximately 85% with surgery.
- Aftercare: Your dog may need rest, restricted activity, and follow-up appointments to monitor healing.

3. Hip Toggle Surgery

In this surgical option, a toggle pin and suture are used to create a new ligament-like structure to hold the hip in place. It's commonly used for more severe or chronic cases where the hip is unstable.

- Success Rate: Generally effective but has a small risk of the hip re-dislocating.
- Risks: The surgical site may become infected, or the toggle may loosen over time.

4. Femoral Head and Neck Osteotomy (FHNO) or Total Hip Replacement (THR)

These are more advanced surgeries and are often considered for dogs with severe damage or hip dysplasia. In an FHNO, the head of the femur is removed to reduce pain, allowing a "false joint" to form. THR replaces the hip joint entirely and is considered the gold standard in these cases especially for larger or active dogs.

VETERINARY SPECIALIST ADVICE

© 2025 Veterinary Specialist Advice (VSA). All rights reserved.Educational use only; VSA and its authors are not responsible for outcomes and this does not replace professional veterinary judgment.

Hip Luxation in Dogs

Post-Treatment Care

- Restricted Movement: After any treatment, it's essential to limit your dog's activity to allow the hip to heal. This may mean crate rest, short walks on a leash, and no jumping.
- Pain Management: Pain relief medication will be prescribed to keep your dog comfortable.
- Physical Therapy: In some cases, gentle exercises or physical therapy can help restore strength and movement.

Potential Complications

- Reluxation (Re-Dislocation): This can occur, especially in non-surgical treatments.
- Infection: Any surgical site can become infected, which may need antibiotics or additional care.
- Nerve Injury: Rarely, nearby nerves may be affected during surgery, potentially leading to temporary weakness.
- Arthritis: Over time, dogs with a history of hip luxation may develop arthritis in the hip, which could cause stiffness and pain as they age.

Outlook

With the right treatment and care, most dogs recover well from hip luxation and can return to normal activities. However, it's important to follow the vet's post-treatment instructions to ensure the best outcome and reduce the risk of future problems.

Hip Toggle Surgery for Dogs

What is Hip Toggle Surgery?

Hip toggle surgery is a commonly performed surgical procedure used to treat hip luxation (dislocation), where the femoral head (top part of the thigh bone) comes out of the hip socket. This technique is especially useful in cases where the hip continues to dislocate after other methods, like closed reduction, have been tried. Hip toggle surgery is designed to create stability in the hip joint by mimicking the ligament that normally holds the femoral head in place.

How Does Hip Toggle Surgery Work?

In hip toggle surgery, the veterinarian places a strong suture (a type of surgical thread) through the bone of the femur and the pelvis, using a small device called a "toggle pin." This pin acts like an anchor within the hip socket, securing the femur in place and creating a new, stable connection between the femur and the hip socket. Essentially, the suture serves as a replacement for the ligament that usually holds the hip joint together.

VETERINARY SPECIALIST ADVICE

© 2025 Veterinary Specialist Advice (VSA). All rights reserved.Educational use only; VSA and its authors are not responsible for outcomes and this does not replace professional veterinary judgment.

Hip Luxation in Dogs

Benefits of Hip Toggle Surgery

- Stability: The suture acts like a new ligament, keeping the femur in place within the hip socket.
- Reduced Risk of Re-Dislocation: This technique provides a long-term solution to prevent the hip from dislocating again.
- Minimally Invasive: Compared to more extensive surgeries, hip toggle surgery is less invasive and can be less painful.

Potential Risks and Complications

Like all surgeries, hip toggle surgery has some potential risks:

- Re-Luxation: There is a small chance that the hip could dislocate again, particularly if the dog is very active during recovery.
- Infection: The surgical site can become infected, requiring additional treatment.
- Suture Wear and Tear: Over time, the suture can wear down, potentially leading to loosening. However, with appropriate activity restrictions and care, this risk is minimized.

Recovery After Hip Toggle Surgery

After the surgery, your dog will need a period of rest and restricted activity to allow the joint to heal properly. Typical post-surgery recommendations include:

- Restricted Exercise: Avoid running, jumping, or rough play for 6-8 weeks.
- Short Walks: Leash walks only for bathroom breaks are allowed during early recovery.
- Medication: Pain relief and anti-inflammatory medications to keep your dog comfortable.
- Follow-Up Visits: Regular check-ups with the vet to ensure the hip is healing correctly.


Outcome

For many dogs, hip toggle surgery provides excellent stability, allowing them to return to a comfortable, active life. While long-term joint health should still be monitored, this procedure typically offers a reliable solution for dogs with hip dislocations.

VETERINARY SPECIALIST ADVICE

© 2025 Veterinary Specialist Advice (VSA). All rights reserved.Educational use only; VSA and its authors are not responsible for outcomes and this does not replace professional veterinary judgment.

Hip Luxation in Dogs



VETERINARY SPECIALIST ADVICE

© 2025 Veterinary Specialist Advice (VSA). All rights reserved.Educational use only; VSA and its authors are not responsible for outcomes and this does not replace professional veterinary judgment.